


Data Sheet

HUMAN UROKINASE PLASMINOGEN ACTIVATOR-RECEPTOR [UPA-R]

ANTIBODY, POLYCLONAL

Catalog no.:	A 7612.1 / A 7612.2
Immunogen:	Purified human Urokinase Plasminogen Activator-Receptor
Synonyms:	Urokinase plasminogen activator surface receptor, U-PAR, uPAR, Monocyte activation antigen Mo3, CD87
Swiss-Prot No:	Q03405
Gene Information:	Gene Name: PLAUR, MO3, UPAR
	GeneID: 5329
Host:	Goat
Matrix:	Serum
Specificity:	Human Urokinase Plasminogen Activator-Receptor
	There were no cross reactivities obtained with human tPA and human PAI-2
Contents:	20 µl / 100 µl (lyophilized)
	Resuspend in 20 µl / 100 µl aqua bidest.
Known applications:	ELISA, Immunohistochemistry (Frozen sections & paraffin sections)
	Recommended procedure for immunohistochemistry:
	Dilutions for frozen sections: 1:50 – 1: 100
	Dilutions for paraffin sections: 1:50
	Recommended procedure for paraffin sections: Pretreatment with 0.05% Pronase E (Sigma) for 20 minutes.
	This antibody has not been tested for use in all applications. This does not necessarily exclude its use in non-tested procedures. The stated dilutions are recommendations only. End users should determine optimal dilutions in their system using appropriate negative/positive controls.
Store at:	2-8 °C (lyophilized); - 20 °C (dissolved)
	Repeated thawing and freezing must be avoided
Last updated on:	13 June 2016

Immundiagnostik AG
Stubenwald-Allee 8a · 64625 Bensheim · Germany
Phone: +49 (0) 62 51/70 19 00 · Fax: +49 (0) 62 51/84 94 30 · info@immundiagnostik.com · www.immundiagnostik.com



For research use only

Publishing research using A 7612? Please let us know so that we can cite your publication as a reference.

Immundiagnostik AG
Stubenwald-Allee 8a · 64625 Bensheim · Germany
Phone: +49 (0) 62 51/70 19 00 · Fax: +49 (0) 62 51/84 94 30 · info@immundiagnostik.com · www.immundiagnostik.com